

ASSESSMENT FOR SUSPECTED COVID-19

LEGEND

EBR: Evidence-Based Recommendation
CBR: Consensus-Based Recommendation
PP: Practice Point

Living
guidance

Not prioritised
for review

VERSION 3.0

PUBLISHED
10 SEPTEMBER 2020

INITIAL SCREENING

GENERAL

Assessment via telehealth

Where possible, undertake initial screening and assessment over the phone or by video. Video is preferred as it provides additional visual cues and therapeutic presence and may prevent the need for an in-person visit. **PP** [BMJ]

Speak to the affected person if possible, rather than their carer or a family member. **PP** [BMJ]

Assessment in-person

If a person with symptoms suggestive of COVID-19 presents to a clinic without a prior telehealth assessment, undertake the initial screening outside the clinic or in an area away from other people. **PP** [Taskforce]

SYMPTOMS AND SIGNS

Ask about the following:

- date of onset of symptoms
- key symptoms: fever, shortness of breath, dry cough, muscle aches, tiredness
- other symptoms: sore throat, headache, runny nose, diarrhoea and nausea.

PP [BMJ]

EPIDEMIOLOGY

For people in the community

Ask about the following in the 14 days prior to illness onset:

- Close contact with a confirmed or probable case of COVID-19
- International or interstate travel
- Passengers and crew who have travelled on a cruise ship
- Healthcare, aged or residential care workers and staff with direct patient contact
- People who have lived in or travelled through a geographically localised area with elevated risk of community transmission, as defined by public health authorities. **PP** [CDNA]

For hospitalised patients

Suspect COVID-19 where no other clinical focus of infection or alternate explanation of the patient's illness is evident. **PP** [CDNA]

Rural and remote access to care

This flowchart should be applied after considering features of the individual patient, their preferences and the context in terms of rurality/remoteness, public health responses and proximity to higher-level care. Application of the flowchart will vary with local current COVID-19 prevalence and availability of testing. Early transfer to a major centre should be considered for those at risk of deterioration.

FOLLOW UP

LABORATORY TESTING

Refer to local testing criteria for SARS-CoV-2, noting that testing criteria may differ between States and Territories. **PP** [Taskforce]

Arrange for both oro- and nasopharyngeal testing (with sputum collection where this can be collected spontaneously and safely). **PP** [Taskforce/CDNA]

Likelihood of COVID-19



Other clinical factors

Potential risk factors for more severe illness
 Older Age
 Indigenous
 History of smoking
 Comorbidities:

- lung disease, including COPD, asthma, or bronchiectasis
- cardiovascular disease, including hypertension
- immunocompromised states (e.g. diabetes, chronic kidney or liver disease, taking chemotherapy, steroids, or other immunosuppressants).

PP [Taskforce/BMJ]

Symptom severity
 Assess the degree of breathlessness by asking the person to describe:

- their presenting problem in their own words, whilst assessing the ease and comfort of their speech.
- the impact of their symptoms on their usual daily activities. Focus on any changes in breathing from normal, such as a new audible wheeze.

PP [BMJ]

Be aware of differential diagnoses

- Serious differential diagnoses include bacterial pneumonia, meningitis, and sepsis
- Influenza is more likely to produce body aches, whilst COVID-19 is more likely to produce shortness of breath.

PP [BMJ]

IN PERSON

Symptoms and signs
 Only follow up with an in-person assessment if a diagnosis of moderate or severe illness cannot be confidently excluded via telehealth assessment or by initial face-to-face screening.
PP [Taskforce]


Managing risk of infection
 Follow national advice for use of PPE in non-inpatient healthcare settings during the COVID-19 outbreak.
PP [Taskforce/AHPPC]

Examination
 Undertake the in-person assessment as per **Other clinical factors** and in addition:

- check temperature, pulse, respiratory rate and effort, degree of cough, and presence of sore throat
- assess oxygen saturation (SaO₂), and if normal, consider repeating after gentle exercise (e.g. walking around the clinic carpark).

PP [Taskforce]

Care pathway



Sources

AHPPC – Australian Health Protection Principal Committee (AHPPC). Guidance on use of personal protective equipment (PPE) in non-inpatient healthcare settings, during the COVID-19 outbreak. 17 June 2020.

BMJ – Covid-19: a remote assessment in primary care. BMJ 2020;368:m1182 doi: 10.1136/bmj.m1182 (25 March 2020)

CDNA – Coronavirus Disease 2019 (COVID-19) Communicable Diseases Network Australia (CDNA) National Guidelines for Public Health Units. V3.8, 23 August 2020

Taskforce – Current guidance from the National COVID-19 Clinical Evidence Taskforce